

Open Guard Edges and Edge Guards in Simple Polygons

Csaba D. Tóth*

Godfried Toussaint†

Andrew Winslow‡

Abstract

An *open edge* of a simple polygon is the set of points in the relative interior of an edge. We revisit several art gallery problems, previously considered for closed edge guards, using open edge guards. A *guard edge* of a polygon is an edge that sees every point inside the polygon. We show that every simple non-starshaped polygon admits at most one open guard edge, and give a simple new proof that it admits at most three closed guard edges. We characterize open guard edges, and derive an algorithm that finds all open guard edges of a simple n -gon in $O(n)$ time in the RAM model of computation. Finally, we present lower bound constructions for simple polygons with n vertices that require $\lfloor n/3 \rfloor$ open edge guards, and conjecture that this bound is tight.

1 Introduction

Let P be a simply connected closed polygonal domain with n vertices. Two points $p, q \in P$ are mutually visible to each other if the closed line segment pq lies in P . In a starshaped polygon P , all points in P are visible from a single point $x \in P$, which is called a *guard point* for P . The set of all guard points is the *kernel* of P .

For a set $S \subseteq P$ of multiple guards, or the trajectories of mobile guards, we adopt the notion of weak visibility [2]. A point $p \in P$ is (weakly) visible to a set $S \subseteq P$ if it is visible from some point in S . If every point $p \in P$ is (weakly) visible from S , then S is a *guard set*.

Park *et al.* [8] considered guard sets restricted to (closed) edges of a polygon. They proved that a non-starshaped simple polygon has at most three closed guard edges, and this bound is tight. They also designed an $O(n)$ time algorithm for finding all closed guard edges in a simple n -gon. Later, it was shown that a shortest guard segment along the boundary of P , or anywhere in P can also be found in optimal $O(n)$ time [3, 4]. A *watchman tour* is a closed curve $\gamma \subset P$ which is a guard set for P . Tan [11] gave an $O(n^5)$ time algorithm for finding a *shortest* watchman tour.

*Department of Mathematics and Statistics, University of Calgary, Calgary, AB, Canada, cdtoth@math.ucalgary.ca

†Department of Music, Harvard University, Cambridge, MA, USA, Department of Computer Science, Tufts University, Medford, MA, USA, School of Computer Science, McGill University, Montreal, QC, Canada, godfried@cs.mcgill.ca

‡Department of Computer Science, Tufts University, Medford, MA, USA awinslow@cs.tufts.edu

If several guards are available, we are interested in the minimum number of guards that can jointly cover any simple polygon with n vertices. By a classical result of Klee, a set of $\lfloor n/3 \rfloor$ vertex guards are always sufficient and sometimes necessary to cover a simple n -gon. It is known that $\lfloor n/4 \rfloor$ closed edge guards are sometimes necessary, and $\lfloor 3n/10 \rfloor + 1$ are always sufficient [9]. It is a longstanding conjecture that $\lfloor n/4 \rfloor + O(1)$ closed edge guards are always sufficient. However, $\lfloor n/4 \rfloor$ (open or closed) segment guards are always sufficient and sometimes necessary [7].




Figure 1: The region visible by an open edge uv (left) and a closed edge uv (right) in a simple polygon.

Viglietta [12] recently suggested the use of open edge guards for various scenarios. A *closed edge* includes the endpoints, and an *open edge* does not. See Fig. 1. Intuitively, a closed edge can “see around the corner” if its endpoint is a reflex vertex, while an open edge cannot. In this note, we examine two art gallery problems involving edges of polygons. First, *guard edges* of a polygon; single edges that guard the entire polygon. Then we consider *edge guards*; sets of edges that together guard the entire polygon.

2 Preliminaries

It is easy to express visibility in terms of shortest paths in a simple polygon (*c.f.*, [1]). Given two points, p and q , inside a simple polygon P , the geodesic path(p, q) is the shortest directed path from p to q that lies entirely in P . Points p and q see each other iff path(p, q) is a straight line segment. Every interior vertex of path(p, q) is a reflex vertex of P . We characterize weak visibility between a point and an edge in terms of geodesics.

Lemma 1 *Let p be a point inside a simple polygon P .*

- Point p is visible from an open edge uv iff p is the only common vertex of path(p, u) and path(p, v);*
- p is visible from a closed edge uv iff all common vertices of path(p, u) and path(p, v) are in $\{p, u, v\}$.*

Proof. (a) If p is the only common vertex of the two geodesics, then uv , $\text{path}(p, u)$, and $\text{path}(p, v)$ form a pseudo-triangle lying in P with corners p, u and v . Each corner of a pseudo-triangle is weakly visible from the opposite side, hence p is visible from a point in uv (Fig. 2, left). If $q \neq p$ is the last common vertex of the two geodesics, then q is an interior point of every geodesic from p to any $w \in uv$, hence p is not visible from any point of the open edge uv (Fig. 2, middle).




Figure 2: The geodesics $\text{path}(p, u)$ and $\text{path}(p, v)$. Left: p is the only common vertex of $\text{path}(p, u)$ and $\text{path}(p, v)$. Middle: the common vertices are p and q . Right: The common vertices are p and v .

(b) If p is the only common vertex of the two geodesics, then p is visible from an interior point of uv as in part (a). If u or v is the only common vertex (apart from p) of the two geodesics, then point p is directly visible from u or v (Fig. 2, right). Finally, if $q \notin \{p, u, v\}$ is a common vertex of the two geodesics, then q is an interior point of every geodesic from p to any $w \in uv$, and hence p is not visible from any point of the closed edge uv . \square

3 Open Guard Edges

In this section we consider open guard edges. Observe that every edge of a convex polygon is a guard edge, since it lies in the kernel of the polygon; but there may be up to $n/4$ open guard edges even if all edges are disjoint from the kernel (Fig. 3, left). In this section, we show that every non-starshaped simple polygon has at most one open guard edge. This bound is tight, as shown by the example in Fig. 3, right.




Figure 3: Left: a starshaped n -gon P with $n/4$ open guard edges where the kernel lies in the interior of P . Right: a non-starshaped polygon with one open guard edge.

We prove the upper bound by contradiction: we prove that a simple polygon with at least two open guard edges is starshaped. Let P be a simple polygon, and

suppose that edges ab and cd are open guard edges. We may assume without loss of generality that a, b, c, d are in counterclockwise order along the boundary of P (possibly, $b = c$ or $d = a$).

Lemma 2 $\text{path}(b, c)$ and $\text{path}(a, d)$ are disjoint. $\text{path}(a, c) = ac$ and $\text{path}(b, d) = bd$ are line segments.

Proof. Note that $ab, \text{path}(b, c), cd,$ and $\text{path}(a, d)$ form a geodesic quadrilateral Q . Every geodesic between a point in ab and a point in cd lies in Q . If $\text{path}(b, c)$ and $\text{path}(a, d)$ have a common interior vertex q , then a or b is not visible from the open edge cd by Lemma 1, and so cd cannot be a guard edge. We conclude that $\text{path}(b, c)$ and $\text{path}(a, d)$ are disjoint, and Q is a simple polygon.

The geodesics $\text{path}(a, c)$ and $\text{path}(b, d)$ lie in Q , so any interior vertex of $\text{path}(a, c)$ and $\text{path}(b, d)$ is a vertex of Q . If an interior vertex of $\text{path}(a, c)$ is in $\text{path}(b, c)$, then c is not visible from ab . Similarly, if an interior vertex of $\text{path}(a, c)$ is in $\text{path}(a, d)$, then a is not visible from cd . Hence, $\text{path}(a, c)$ has no interior vertices. Analogous argument shows that $\text{path}(b, d)$ has no interior vertices, either. \square

Lemma 3 The intersection point $x = ac \cap bd$ is in the kernel of P .

Proof. Refer to Fig. 4. It is enough to show that an




Figure 4: A schematic of the proof that a simple polygon with two open guard edges must be starshaped. The guard edges are ab and cd . The point $x = ac \cap bd$ is in the kernel of the polygon, since every point $p \in P$ is visible from x .

arbitrary point p in polygon P is visible from x . By Lemma 2, ac and bd are diagonals of P . The triangles $\Delta(abx)$ and $\Delta(cdx)$ lie inside P . If $p \in \Delta(abx)$ or $p \in \Delta(cdx)$, then segment px lies in the same triangle. Assume now that p is outside of both triangles. Since ab and cd are open guard edges, p sees some points in their relative interiors, say $o \in ab$ and $q \in cd$. The quadrilateral $Q = (o, p, q, x)$ is simple, and so it lies inside P . Note that Q has convex vertices at o and q . No matter whether Q is a convex or a non-convex quadrilateral, its diagonal px lies inside Q , and hence inside P . \square

Theorem 4 Every non-starshaped simple polygon has at most one open guard edge.

Proof. If a simple polygon has two open guard edges, then it has a nonempty kernel by Lemma 3, and thus is starshaped. So every non-starshaped simple polygon has at most one open guard edge. \square

Remark. The upper bound of Theorem 4 does not apply to polygons with holes. Note that an open edge on the boundary of a hole cannot see the entire boundary of the hole. So all open edge guards are on the boundary of the outer polygon. By the result in [8] there are at most 3 closed guard edges on the outer boundary of a polygon with holes. Since every open guard edge is a closed guard edge, as well, a polygon with holes has at most 3 open guard edges. This upper bound is tight, as shown by the following simple construction. Let the outer polygon and a hole be two centrally dilated triangles. Then all three open edges of the outer polygon are guard edges.

4 Closed Guard Edges

In this section, we extend the argument of the previous section to give a short proof for the following result of Park *et al.* [8].

Theorem 5 ([8]) *Every non-starshaped simple polygon has at most three closed guard edges.*

We proceed by contradiction, and show that the presence of four closed guard edges implies that the polygon is starshaped. Let P be a simple polygon where $g_1, g_2, g_3,$ and $g_4,$ in counterclockwise order, are guard edges. Let $g_1 = ab$ and $g_3 = cd$ such that $a, b, c,$ and d are in counterclockwise order along P . Note that the vertices $a, b, c,$ and d are distinct.

Lemma 6 *The geodesics $\text{path}(b, c)$ and $\text{path}(a, d)$ are disjoint; and all vertices of the geodesics $\text{path}(a, c)$ and $\text{path}(b, d)$ are in $\{a, b, c, d\}$.*

Proof. Consider the geodesic quadrilateral Q formed by $ab, \text{path}(b, c), cd,$ and $\text{path}(a, d)$. Every geodesic between a point in ab and a point in cd lies in Q . Suppose that an interior vertex q of $\text{path}(b, c)$ is a vertex of $\text{path}(a, d)$. If $q = a$ or an interior vertex of $\text{path}(a, d)$, then b is not visible from the closed edge cd by Lemma 1. Similarly, if $q = d$, then c is not visible from the closed edge ab . We conclude that $\text{path}(b, c)$ and $\text{path}(d, a)$ are disjoint, and Q is a simple polygon.

The geodesics $\text{path}(a, c)$ and $\text{path}(b, d)$ lie in Q , so any interior vertex of $\text{path}(a, c)$ and $\text{path}(b, d)$ is a vertex of Q . If $\text{path}(a, c)$ and $\text{path}(b, c)$ have a common interior vertex, then c is not visible from ab . Similarly, no two geodesics from $\{a, b\}$ to $\{c, d\}$ can have any common interior vertex. Hence all interior vertices of $\text{path}(a, c)$ and $\text{path}(b, d)$ are in $\{a, b, c, d\}$. \square

Corollary 7

- If $\{a, b, c, d\}$ is in convex position, then $\text{path}(a, c) = ac$ and $\text{path}(b, d) = bd$. Fig. 5, left.
- Otherwise suppose w.l.o.g. that $\text{conv}(\{a, b, c, d\}) = \Delta(abc)$. Then $\text{path}(a, c) = (a, d, c)$ and $\text{path}(b, d) = bd$. Fig. 5, right.




Figure 5: The convex hull of two closed guard edges, ab and cd , is either a quadrilateral or a triangle.

Lemma 8 *The intersection point $x = \text{path}(a, c) \cap \text{path}(b, d)$ is in the kernel of P .*

Proof. It is enough to show that an arbitrary point p in polygon P is visible from x . By Corollary 7, the triangles $\Delta(abx)$ and $\Delta(cdx)$ lie inside P (one of the triangles may be degenerate). If p is in $\Delta(abx)$ or $\Delta(cdx)$, then segment px lies in the same triangle. Refer to Fig. 6.




Figure 6: A schematic of the proof that a simple polygon with four closed guard edges must be starshaped. Suppose that $g_1 = ab, g_2, g_3 = cd,$ and g_4 are guard edges. If a point $p \in P$ is not visible from $x = \text{path}(a, c) \cap \text{path}(b, d)$, then we show that p is also not visible from g_2 or g_4 .

Assume now that p is outside of both triangles and, w.l.o.g. it is on the right side of the directed geodesics $\text{path}(a, c)$ and $\text{path}(b, d)$. That is, p and the guard edge g_4 are on opposite sides of these geodesics.

If $\text{path}(p, x) = px$, then p is visible from x . Suppose, to the contrary, that $\text{path}(p, x)$ is not a straight line segment. Assume w.l.o.g. that $\text{path}(p, x)$ makes a right turn at its last interior vertex q . Then $\text{path}(p, d)$ also makes a right turn at q . Since p is visible from the guard edge cd , we must have $q = c$ by Lemma 1(b). Recall that any geodesic from p to a point in g_4 crosses both $\text{path}(a, c)$ and $\text{path}(b, d)$. Since $\text{path}(p, x)$ makes a right turn at c , every geodesic from p to a point in

g_4 also make a right turn at c . However, c is disjoint from g_4 , and by Lemma 1(b), p is not visible from g_4 , contradicting our initial assumption. We conclude that $\text{path}(p, x)$ is a straight line segment, and so p is visible from x . \square

Proof of Theorem 5. If a simple polygon has four closed guard edges, then it has a nonempty kernel by Lemma 8, and thus is starshaped. So every non-starshaped simple polygon has at most three closed guard edges. \square

5 Characterizing Open Guard Edges

In this section, we characterize the open guard edges of a simple polygon P in terms of the left and right kernels of P (defined below). This leads to a straightforward algorithm to find all open guard edges in P .

Left and right kernels. Recall that the set of points from which the entire polygon P is visible is the *kernel*, denoted $K(P)$, which is the intersection of all halfplanes bounded by a supporting line of an edge of P and facing towards the interior of P . Lee and Preparata [5] designed an optimal $O(n)$ time algorithm for computing the kernel of simple polygon with n vertices. We now define a weaker version of the kernel: the *left* and *right kernels* of P , denoted $K_{\text{left}}(P)$ and $K_{\text{right}}(P)$.

For every reflex vertex r , we define two polygons $C_{\text{left}}(r) \subset P$ and $C_{\text{right}}(r) \subset P$. Shoot a ray from r in a direction collinear with the edge incident to r preceding (resp., following) r in counterclockwise order; and let $C_{\text{left}}(r)$ (resp., $C_{\text{right}}(r)$) be the part of P on the left (resp., right) of the ray. These polygons have previously been defined in [3]. It is clear that if P is weakly visible from a set $S \subset P$, then S must intersect both $C_{\text{left}}(r)$ and $C_{\text{right}}(r)$ for every reflex vertex r .

Now we define $K_{\text{left}}(P)$ as the intersection of polygons $C_{\text{left}}(r)$ for all reflex vertices r ; and $K_{\text{right}}(P)$ as the intersection of polygons $C_{\text{right}}(r)$ for all r . See Fig. 7 for an example. Clearly, we have

$$K(P) = K_{\text{left}}(P) \cap K_{\text{right}}(P).$$

By construction, both $K_{\text{left}}(P)$ and $K_{\text{right}}(P)$ are convex polygons, whose edges are collinear with some of the edges of P .

Left and right kernel decompositions. In the following lemma we use two decompositions of a polygon based on its left and right kernels. The *left kernel decomposition* is the decomposition of the polygonal domain P in the following way: One cell of the decomposition is the left kernel $K_{\text{left}}(P)$. The region inside P but in the exterior of $K_{\text{left}}(P)$ is decomposed by extending each edge of $K_{\text{left}}(P)$ in clockwise direction. Refer to Fig. 7. Since $K_{\text{left}}(P)$ lies on the left side of rays




Figure 7: The left and right kernels of a polygon. The dotted lines bound some polygons $C_{\text{left}}(r)$ and $C_{\text{right}}(r)$, but they are not part of the kernel decompositions.

emitted from reflex vertices of P , the clockwise extension of every edge of $K_{\text{left}}(P)$ reaches a collinear edge of P . The right kernel decomposition is defined analogously: one cell is $K_{\text{right}}(P)$, and the rest of P is decomposed by counter-clockwise extensions of the edges of $K_{\text{right}}(P)$. Note that if an open edge of P is disjoint from the left kernel, then it is adjacent to a unique region of the left kernel decomposition. Additionally, each region of the decomposition, except for $K_{\text{left}}(P)$, has exactly one common edge with the left kernel.

Lemma 9 *An open edge e of a simple polygon P is a guard edge of P iff e intersects both the left and the right kernels of P .*

Proof. Let $e = uv$ be an open edge of P . First assume that e is disjoint from the left kernel $K_{\text{left}}(P)$. Then e is adjacent to a unique region in the left kernel decomposition of P . This region is adjacent to a unique edge k of $K_{\text{left}}(P)$, and k lies on a ray emitted by a reflex vertex r on P . Then e and the polygon $C_{\text{left}}(r)$ lies on opposite sides of the supporting line of k . Hence e does not intersect $C_{\text{left}}(r)$, and so it is not a guard edge.

Now assume that $e = uv$ is not a guard edge, that is, there is a point $p \in P$ such that p is not visible from e . By Lemma 1(a), the geodesics $\text{path}(p, u)$ and $\text{path}(p, v)$ have common interior vertices. Let r be their last common vertex, which is necessarily a reflex vertex of P , and assume w.l.o.g. that both geodesics make a right turn at r . Then $p \in C_{\text{left}}(r)$, but e is disjoint from $C_{\text{left}}(r)$. That is, e is disjoint from the left kernel of P . \square

Finding all open guard edges in $O(n)$ time. We use Lemma 9 to create a simple $O(n)$ time algorithm for finding all open guard edges of a simple polygon P with n vertices (independent of whether P is starshaped or not). The left and right kernels of P can be computed in $O(n)$ time using a modified version of the algorithm of Lee and Preparata [5], originally designed for computing the kernel $K(P)$. For each edge of the left and right kernels, mark any intersection with the collinear edge of P . Now check the marks on all edges of P in $O(n)$

time, and report those that intersect both the left and the right kernels.

6 Open Edge Guards

Recall that every simple polygon with n vertices can be covered by $\lfloor 3n/10 \rfloor + 1$ closed edge guards, and there are n -gons that require at least $\lfloor n/4 \rfloor$ closed edge guards. It turns out that the endpoints of the closed edge guards are crucial for these bounds. Significantly more edge guards may be necessary if the endpoints are removed.

We construct four different infinite families of polygons that require $\lfloor n/3 \rfloor$ open edge guards for n vertices. Refer to Fig. 8. The lower bounds for all our constructions can be verified by a standard “hidden point” argument. We hide $\lfloor n/3 \rfloor$ points (gray dots in Fig. 8) in the interior of a polygon such that each open edge guard sees exactly one such point. That is, each hidden point requires a unique open edge guard, and any set of fewer than $\lfloor n/3 \rfloor$ open edge guards would miss at least one hidden point.

It is not difficult to see that $\lfloor n/2 \rfloor$ open edge guards are always sufficient. Partition the set of edges of the polygon into two subsets for which the interior normals of the edges have either a positive or negative y -component. Each subset of open edges jointly covers the entire polygon. We conjecture this upper bound is weak, and that $\lfloor n/3 \rfloor$ is the tight bound for the number of open edge guards necessary and sufficient to guard any simple polygon with n vertices.

References

- [1] D. Avis, T. Gum, and G. T. Toussaint, Visibility between two edges of a simple polygon, *The Visual Computer*, 2:342–357, 1986.
- [2] D. Avis and G. Toussaint, An optimal algorithm for determining the visibility of a polygon from an edge, *IEEE Tran. Comput.* C-30:910–914, 1981.
- [3] B. K. Bhattacharya, G. Das, A. Mukhopadhyay, and G. Narasimhan, Optimally computing a shortest weakly visible line segment inside a simple polygon, *Comput. Geom. Theory. Appl.*, 23:1–29, 2002.
- [4] D. Z. Chen, Optimally computing the shortest weakly visible subedge of a simple polygon, *J. Algorithms* 20:459–478, 1996.
- [5] D. T. Lee and F. Preparata, An optimal algorithm for finding the kernel of a polygon, *J. ACM* 26:415–421, 1979.
- [6] B.-K. Lu, F.-R. Hsu, and C. Y. Tang, Finding the shortest boundary guard of a simple polygon, *Theor. Comp. Sci.*, 263:113–121, 2001.
- [7] J. O’Rourke, Galleries need fewer mobile guards: A variation on Chvatal’s theorem, *Geometriae Dedicata* 14:273–283, 1983.
- [8] J. Park, S. Y. Shin, K. Chwa, and T. C. Woo, On the number of guard edges of a polygon, *Discrete Comput. Geom.*, 10:447–462, 1993.
- [9] T. C. Shermer, Recent results in art galleries, *Proc. IEEE*, 80:1384–1399, 1992.
- [10] S. Y. Shin and T. Woo, An optimal algorithm for finding all visible edges in a simple polygon, *IEEE Tran. Robotics and Automation* 5:202–207, 1989.
- [11] X. Tan, Fast computation of shortest watchman routes in simple polygons, *Inf. Proc. Lett.* 77:27–33, 2001.
- [12] G. Viglietta, Searching polyhedra by rotating planes, manuscript, arXiv:1104.4137, 2011.




Figure 8: Examples of polygons requiring $n/3$ open edge guards. The gray dots in each polygon indicates a set of points that require a distinct edge to guard each point.